



Cholesterol modulates bitter taste receptor function

Sai Prasad Pydi^{a,b}, Md. Jafurulla^c, Lisa Wai^a, Rajinder P. Bhullar^a,
Prashen Chelikani^{a,b,*}, Amitabha Chattopadhyay^{c,**}

^a Department of Oral Biology and Manitoba Chemosensory Biology Research Group, University of Manitoba, Winnipeg, MB R3E 0W4, Canada

^b Children's Hospital Research Institute of Manitoba (CHIRM), Winnipeg, MB R3E 0W4, Canada

^c CSIR-Centre for Cellular and Molecular Biology, Uppal Road, Hyderabad 500 007, India

ARTICLE INFO

Article history:

Received 19 November 2015

Received in revised form 2 June 2016

Accepted 7 June 2016

Available online 08 June 2016

Keywords:

Bitter taste receptors

GPCR

T2R4

Cholesterol

CRAC

Calcium signaling

ABSTRACT

Bitter taste perception in humans is believed to act as a defense mechanism against ingestion of potential toxic substances. Bitter taste is perceived by 25 distinct bitter taste receptors (T2Rs) which belong to the family of G protein-coupled receptors (GPCRs). In the overall context of the role of membrane lipids in GPCR function, we show here that T2R4, a representative member of the bitter taste receptor family, displays cholesterol sensitivity in its signaling function. In order to gain further insight into cholesterol sensitivity of T2R4, we mutated two residues Tyr114^{3.59} and Lys117^{3.62} present in the cholesterol recognition amino acid consensus (CRAC) motif in T2R4 with alanines. We carried out functional characterization of the mutants by calcium mobilization, followed by cholesterol depletion and replenishment. CRAC motifs in GPCRs have previously been implicated in preferential cholesterol association. Our analysis shows that the CRAC motif represents an intrinsic feature of bitter taste receptors and is conserved in 22 out of 25 human T2Rs. We further demonstrate that Lys117, an important CRAC residue, is crucial in the reported cholesterol sensitivity of T2R4. Interestingly, cholesterol sensitivity of T2R4 was observed at quinine concentrations in the lower mM range. To the best of our knowledge, our results represent the first report addressing the molecular basis of cholesterol sensitivity in the function of taste receptors.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

G protein-coupled receptors (GPCRs) are cellular nanomachines that represent the most important class of membrane-embedded receptors in eukaryotes. They act as signaling hubs and mediate diverse cellular responses [1,2]. As a consequence, GPCRs constitute the largest family of clinical drug targets [3]. The human bitter taste receptors (T2Rs) are members of GPCR family. They display similarities and differences with Class A GPCRs [4,5]. T2Rs are chemosensory receptors with important therapeutic potential [6,7]. Bitter taste perception is believed to act as a defense mechanism against ingestion of potential toxic substances. In humans, bitter taste is perceived by 25 distinct T2Rs. They are expressed in the oral cavity, gastrointestinal neuroendocrine cells of the large intestine and in many extraoral tissues [8,9]. T2Rs typically consist of 290–333 amino acids and display considerable degree of evolutionary divergence [10].

The role of membrane lipids in GPCR function is a promising and emerging area of research [11–17]. In this context, membrane cholesterol enjoys a unique position in modulating GPCR organization, dynamics and function [11–16,18]. Most of the work involving GPCR-cholesterol interaction has been carried out with Class A GPCRs, mainly the serotonin_{1A} receptor, β_2 -adrenergic receptor and rhodopsin. Work from a number of laboratories has shown that the function of these receptors depends on membrane cholesterol content, although the exact mechanism for such cholesterol dependence of receptor function is not clear. The effect of cholesterol on GPCR function and organization is believed to be due to direct and/or indirect effects [19]. On the other hand, it has been shown that membrane cholesterol is not necessary for the function of the neurotensin receptor 1, another member of Class A GPCR family [20]. Clearly, it is still early days for predicting whether the function of a specific GPCR would depend on membrane cholesterol. Keeping in mind this overall context, in this work, we have explored the membrane cholesterol sensitivity of a representative bitter taste receptor, T2R4. A schematic representation of the membrane embedded T2R4 is shown in Fig. 1. Our results show that the signaling of T2R4 is sensitive to membrane cholesterol content. In addition, we have analyzed the basis of the cholesterol sensitivity of receptor function in terms of cholesterol recognition/interaction amino acid consensus (CRAC) motif found in T2R4. In T2R4, Leu110^{3.55}, Tyr114^{3.59} and Lys117^{3.62} are the conserved amino acids of the CRAC motif

Abbreviations: CRAC, cholesterol recognition/interaction amino acid consensus; GPCR, G protein-coupled receptor; M β CD, methyl- β -cyclodextrin; RFU, relative fluorescence unit; T2R, bitter taste receptor.

* Corresponding author. Tel.: +1 204 789 3539; fax: +1 204 789 3913.

** Corresponding author. Tel.: +91 40 2719 2578; fax: +91 40 2716 0311.

E-mail addresses: Prashen.Chelikani@umanitoba.ca (P. Chelikani), amit@ccmb.res.in (A. Chattopadhyay).




Fig. 1. A schematic representation of the membrane embedded bitter taste (T2R4) receptor showing its topological and other features. The membrane is shown as a bilayer of phospholipids and cholesterol, representative of typical eukaryotic membranes. The coding region of T2R4 receptor consists of 299 amino acids. The transmembrane helices (I–VII) of the receptor, predicted using TMpred and HMMTOP programs, are depicted as α -helices. The exact boundary between the membrane and the aqueous phase is not known and therefore the locations of the residues relative to the membrane bilayer are putative. The amino acids in the receptor sequence are shown as circles. An octapeptide FLAG sequence was introduced at the N-terminus of the receptor (residues of FLAG sequence are shown as broken circles). The amino acid residues of CRAC motif (see later) are highlighted in yellow and the conserved residues L110, Y114 and K117 are highlighted in red. Adapted and modified from [22]. See text for other details.

(Ballesteros-Weinstein numbering) [5]. Our recent studies showed that the L110A mutation does not affect the expression or function of the receptor [7]. In this study, we mutated the other two CRAC motif residues, Tyr114^{3,59} and Lys117^{3,62}, to alanine in order to understand their role in cholesterol binding and function of T2R4. To the best of our knowledge, this is the first report of cholesterol sensitivity of taste receptor function. Given the importance of cholesterol in human diet, cholesterol sensitivity of bitter taste receptors assumes physiological relevance.

2. Materials and methods

2.1. Materials

Cholesterol, methyl- β -cyclodextrin (M β CD), fetal calf serum, quinine and other chemicals were purchased from Sigma Chemical Co. (St. Louis, MO). Calcium sensitive dye Fluo-4 NW, lipofectamine 2000, D-MEM/F-12 [Dulbecco's modified Eagle medium: nutrient mixture F-12 (Ham) (1:1)], and Amplex Red cholesterol assay kit were purchased from Invitrogen Life Technologies (Carlsbad, CA). All other chemicals used were of the highest purity available.

2.2. Methods

2.2.1. Cell culture and transient transfection

CHO cells were maintained in D-MEM/F-12 (1:1) supplemented with 1.2 g/l of sodium bicarbonate, 10% fetal calf serum, 100 U/ml penicillin and 100 μ g/ml of streptomycin in a humidified atmosphere with 5% CO₂ at 37 °C. Fusion of mammalian expression vectors pcDNA3.1 (Invitrogen; Carlsbad, CA) with the N-terminal FLAG tagged wild type T2R4 and the mutant Y114A and K117A receptor genes were carried out as described earlier [21–23]. CHO cells were transiently transfected with respective plasmid constructs using lipofectamine 2000 reagent as per the manufacturer's instructions.

2.2.2. Cholesterol depletion of cells in culture

Depletion of membrane cholesterol from the cells in culture was carried out using M β CD as described earlier [24] with some modifications. CHO cells transfected with wild type, Y114A or K117A receptor genes in pcDNA3.1 were plated at a density of $\sim 1 \times 10^5$ in clear bottom black

walled 96-well plates in D-MEM/F-12 medium 6–8 h after transfection. Cells were grown for 14–16 h, followed by incubation in serum-free medium for 3 h. Cholesterol depletion was carried out by treating cells with 5 mM M β CD for 30 min at 37 °C [24]. After a wash with serum-free medium, cholesterol-depleted cells were used either for cholesterol replenishment or for calcium mobilization measurements.

2.2.3. Cholesterol replenishment in cholesterol-depleted cells

Cholesterol replenishment in cholesterol-depleted cells was carried out using water-soluble cholesterol-M β CD complex as described earlier [25]. Briefly, cholesterol-depleted CHO cells were replenished with cholesterol by incubating with cholesterol-M β CD complex for 10 min in a humidified atmosphere with 5% CO₂ at 37 °C. The complex was prepared by dissolving the required amount of cholesterol and M β CD in a ratio of 1:10 (mol/mol) in sterile water by constant shaking at room temperature (~ 23 °C). Stock solutions (typically 2:20 mM of cholesterol:M β CD) of this complex were freshly prepared and were added to 2 \times serum-free medium to yield final solution containing 1:10 mM of cholesterol:M β CD complex [25]. After a wash with serum-free medium, these cells were used for calcium mobilization measurements.

2.2.4. Receptor expression and calcium mobilization assays

Cell surface receptor expression was analyzed by ELISA using anti-FLAG antibody. The mutants Y114A and K117A were properly expressed on the cell surface with expression levels of $110 \pm 12\%$ and $80 \pm 5\%$, respectively relative to that of wild type T2R4 (taken as 100%). Calcium mobilization measurements were carried out as described earlier [21,26]. Briefly, calcium sensitive dye Fluo-4 NW was loaded into cells by incubating at 37 °C for 45 min followed by 45 min at ~ 22 °C. Receptor activation was determined by measuring changes in intracellular calcium (monitored by measuring increase in fluorescence) upon stimulation with increasing concentration of quinine (0.078 to 5.0 mM) or buffer alone (for measuring basal activity) using Flexstation-3 fluorescence plate reader (Molecular Devices; Sunnyvale, CA) at 525 nm following excitation at 494 nm. Calcium mobilized was expressed as change in relative fluorescence unit (Δ RFU) after subtracting the responses of cells transfected with the empty vector.

2.2.5. Cell membrane preparation and cholesterol quantification

Cell membranes were prepared from untreated, cholesterol-depleted, and cholesterol-replenished cells as previously described [27]. Briefly, cells were harvested by treatment with 10 mM Tris, 5 mM EDTA, pH 7.4 buffer and homogenized with a Polytron homogenizer. The cell lysate was centrifuged at $500 \times g$ for 10 min and the resulting post-nuclear supernatant was centrifuged at $40,000 \times g$ for 30 min. The pellet was suspended in 50 mM Tris, pH 7.4 buffer. The amount of cholesterol present in these membranes was quantified using Amplex Red cholesterol assay kit [28].

2.2.6. Amino acid sequence analysis and identification of CRAC motif in human T2Rs

All the amino acid sequences of human T2Rs were retrieved from the NCBI database. Multiple sequence alignment was performed using ClustalW multiple sequence alignment program, as described previously [7]. The CRAC motif in human T2Rs (with the conserved tyrosine (Y) along with leucine (L)/valine (V) toward its amino terminus and lysine (K)/arginine (R) toward carboxy terminus, within five residues on either side) were manually identified.

2.2.7. Molecular modeling and docking studies

Three-dimensional molecular model of T2R4 was built using Sybyl-X 2.0 molecular modeling suite (Tripos Inc., USA). Rhodopsin crystal structure (PDB ID:3DQB) was used as a template. First stage minimization was performed using the steepest descent and conjugate gradient algorithms. MD simulation for 10 ns was carried out with a time-step of 2 fs, collecting trajectory data every 500 ps. The SHAKE algorithm, which constraints the hydrogen-heavy atom bonds was applied. Simulations were carried out using Sybyl-X 2.0 at constant temperature (300 K), and pressure (1 atm). Periodic boundary conditions were used to treat long-term electrostatics. T2R4 model obtained this way was used for cholesterol docking. Cholesterol was docked to the identified CRAC motif of T2R4 using the Surflex-Dock docking program within Sybyl-X 2.0. This complex was further energy minimized. PyMol molecular visualizer software was used to analyze the T2R4-cholesterol complex.

2.2.8. Analysis of data

Nonlinear curve fitting of the dose response data was carried out using Graphpad Prism version 4.0 (San Diego, CA). EC_{50} values were calculated from nonlinear regression analysis.

3. Results

3.1. Membrane cholesterol sensitivity of human T2R4

In order to examine the sensitivity of T2R4 function to membrane cholesterol, we monitored the phospholipase C mediated increase in calcium mobilization upon activation of T2R4 with increasing concentration of the specific agonist quinine, under conditions of varying cholesterol content. Quinine is the only pharmacologically characterized T2R4 full agonist, including its ligand binding site mapped at the receptor. We modulated cell membrane cholesterol levels using M β CD, a water-soluble compound that has earlier been shown to selectively and efficiently extract cholesterol from cellular membranes by including it in a central nonpolar cavity [24,29,30]. Fig. 2a shows cell membrane cholesterol content of cholesterol-depleted and cholesterol-replenished cells. Upon treatment of cells with 5 mM M β CD, cholesterol content was reduced to ~66% of control (i.e., without M β CD treatment). Cholesterol replenishment of cholesterol-depleted cells was carried out by incubating the cholesterol-depleted cells with cholesterol-M β CD complex (see Materials and methods). As shown in Fig. 2a, replenishment of cholesterol with this complex resulted in recovery of cholesterol to ~99% of control.

The functional correlates of these manipulations of membrane cholesterol are shown in Fig. 2b. The figure shows a characteristic dose




Fig. 2. Cholesterol-dependent T2R4 signaling. (a) Membrane cholesterol content of CHO cells transfected with T2R4 upon cholesterol depletion and replenishment. Values are expressed as nmol of cholesterol per mg protein. (b) Agonist-stimulated calcium signaling of wild type receptors as a function of membrane cholesterol. Calcium signaling in control (●, olive), cholesterol-depleted (▲, blue), and cholesterol-replenished (■, maroon) CHO cells expressing wild type receptors is shown upon stimulation with increasing concentrations of T2R4 agonist (quinine). CHO cells expressing wild type receptors were treated with M β CD to deplete membrane cholesterol and subsequently treated with cholesterol-M β CD complex to replenish membrane cholesterol. Cells transfected with T2R4 (or the empty vector, pcDNA) were preloaded with the calcium-sensitive dye Fluo-4 NW were stimulated with increasing concentrations of quinine and the resultant calcium mobilization was measured. The difference in relative fluorescence units (RFU) of T2R4 expressing cells with cells transfected with pcDNA is shown as Δ RFU. Nonlinear curve fitting of the dose-response data was carried out using Graphpad Prism version 4.0 (San Diego, CA). Data shown are means \pm S.E. of at least three independent experiments (**corresponds to significant ($p < 0.01$) difference in cholesterol content of cholesterol-depleted cells relative to control cells). See Materials and methods for other details.

response curve for T2R4 in control cells corresponding to EC_{50} value of 1.6 mM (see Table 1). Interestingly, T2R4 was found to exhibit increased signaling upon depletion of membrane cholesterol using M β CD, as evident from the agonist dose response curve. We were not able to determine EC_{50} in cholesterol-depleted cells since the dose response curve did not display saturation at 5 mM quinine concentration. We could not measure calcium responses at quinine concentrations higher than 5 mM due to significant non-specific effects observed at this concentration, as described earlier [22]. Analysis of raw calcium traces, suggest that at 5 mM quinine concentration, the addition of quinine causes an immediate increase in calcium spike (see Fig. S2) and is sustained for an extended period of time. In contrast, at lower concentrations of quinine the initial calcium spike decreases rapidly and approaches baseline. The observed effect at higher quinine concentration could be due to lack of desensitization of T2R4. Fig. 2b shows that agonist mediated calcium signaling was restored similar to control levels (EC_{50} value of 1.8 mM, see Table 1) when cholesterol was replenished close to normal levels. Such characteristic dependence of signaling function on cholesterol content has not been previously demonstrated for any taste receptors.

Table 1

EC₅₀ values for calcium signaling of T2R4 receptors as a function of membrane cholesterol^{a,b}.

	Wild type	Y114A	K117A
Normal	1.6 ± 0.3	1.9 ± 0.3	–
Cholesterol-depleted cells	–	–	–
Cholesterol-replenished cells	1.8 ± 0.5	1.8 ± 0.6	–

^a EC₅₀ values (in mM) were calculated from nonlinear regression analysis of the dose-response data using Graphpad Prism version 4.0.

^b Calcium signaling of the receptor was assayed by monitoring changes in calcium mobilization signal upon stimulation with increasing concentrations of quinine. Data represent means ± S.E. of at least three independent experiments. See [Materials and methods](#) and [Figs. 2–4](#) for more details.

3.2. Cholesterol binding motif in T2Rs

An increasingly emerging feature in recently solved high resolution crystal structures of GPCRs is the presence of closely associated cholesterol molecules ([31–34]; recently reviewed in [15]). Several structural motifs of proteins and receptors have been postulated to induce preferential association with cholesterol [13,15]. A major motif in this category is the CRAC (cholesterol recognition/interaction amino acid consensus) motif [35,36]. The CRAC motif is characterized by the sequence -L/V-(X)_{1–5}-Y-(X)_{1–5}-R/K-, in which (X)_{1–5} represents between one and five residues of any amino acid. Importantly, we recently reported the presence of CRAC motifs in representative GPCRs such as rhodopsin, the β₂-adrenergic receptor, and the serotonin_{1A} receptor [37], that have been shown to exhibit membrane cholesterol sensitivity for their function. High occupancy of cholesterol at some of the CRAC sites in the serotonin_{1A} receptor has been reported by molecular dynamics simulations using the MARTINI coarse-grain approach [38].

In order to explore molecular details of cholesterol sensitivity of T2R4 function (Fig. 2b), we examined whether bitter taste receptors in general, and T2R4 in particular, possess CRAC sequence. We found that the interface of transmembrane helix III and intracellular loop 2 of T2R4 has a CRAC motif (see Fig. 1). Amino acid sequence alignment of human T2Rs showed that the CRAC motif is conserved in 22 out of 25 human T2Rs (see Fig. S1). This implies that CRAC motif is an inherent characteristic feature of majority of T2Rs. Interestingly, we have previously shown that CRAC motifs are inherent characteristic features of the serotonin_{1A} receptor and are conserved over natural evolution [37].

3.3. CRAC motif in T2R4: importance of crucial residues in signaling

In order to understand the importance of conserved CRAC motif residues in T2R4 function, we mutated specific amino acid residues in the CRAC sequence and monitored agonist-stimulated calcium mobilization by the wild type and mutant receptors. In this context, it is important to note that a cholesterol binding motif has previously been proposed to contain an aromatic amino acid (for a possible interaction with ring D of cholesterol) [32] and a positively charged residue (for electrostatic interactions with 3β-hydroxyl group of cholesterol) [39,40]. The crucial CRAC motif residues for T2R4 are leucine, tyrosine, and lysine, at positions 110, 114, and 117, respectively (see Fig. 1). The importance of some of these residues in T2R4 signaling is shown in Fig. 3. The figure shows agonist-stimulated calcium mobilization for the wild type T2R4, and mutant Y114A and K117A receptors with increasing concentration of the T2R4 agonist quinine. Analysis of dose response curves yielded EC₅₀ of 1.6 mM in case of the wild type receptors (see Table 1). The corresponding value for the mutant Y114A receptor was 1.9 mM. Interestingly, the mutant K117A receptor showed increased signaling and did not exhibit saturation even at higher concentrations of quinine. Determination of EC₅₀ was therefore not possible in this case. These results demonstrate that Lys117 in the CRAC sequence has significant effect

on T2R4 signaling with Tyr114 showing only a mild effect. It should be noted here that we have previously shown that the Leu³⁵⁵ (which corresponds to L110) mutation does not affect the expression and function in a similar bitter taste receptor, T2R1 [7].

3.4. Cholesterol sensitivity of mutant receptors: importance of key CRAC residues

The significance of the key amino acid residues of the CRAC sequence could be further explored by monitoring the sensitivity of the function of the mutant receptors under conditions of varying cholesterol content. We therefore modulated membrane cholesterol content of cells expressing the mutant Y114A and K117A receptors and monitored agonist-stimulated calcium signaling for these receptors. Fig. 4a shows that membrane cholesterol content of cells expressing Y114A and K117A receptors was reduced to ~68% and ~38% of control levels upon treatment with MβCD. Cholesterol replenishment of cholesterol-depleted cells using cholesterol-MβCD complex resulted in recovery of cholesterol to ~90% and ~108% of control, respectively. The effect of such modulation of membrane cholesterol content on the signaling of Y114A and K117A receptors is shown in Fig. 4b and c. Fig. 4b shows that the mutant Y114A receptor exhibits concentration-dependent quinine-stimulated calcium signaling with EC₅₀ value of 1.9 mM under control conditions (in the absence of cholesterol modulation), similar to wild type receptor (see Fig. 3 and Table 1). Interestingly, upon depletion of membrane cholesterol, Y114A receptor was found to exhibit enhanced signaling, as evident from the agonist dose response curve (see Fig. 4b). Since the dose response curve did not display saturation, we could not determine EC₅₀ in cholesterol-depleted cells. The figure also shows that agonist mediated calcium signaling was restored to control levels (EC₅₀ value ~1.8 mM, see Table 1) when cholesterol was replenished in cholesterol-depleted cells. On the other hand, the mutant K117A receptor was found to exhibit enhanced signaling relative to the wild type receptor (see Fig. 3 and Table 1). Importantly, the K117A receptor did not exhibit any appreciable change in its signaling upon modulation of membrane cholesterol content (Fig. 4c). Representative raw calcium traces for CHO cells mock-transfected or transfected with Y114A and K117A mutants upon stimulation with quinine under control, cholesterol-depleted and cholesterol-replenished conditions are shown in Fig. S3. Taken together, these results point out that the Lys117 residue of CRAC motif is not only important in normal functioning of the T2R4 receptor but also plays a significant role in sensitivity of the receptor to membrane cholesterol content.




Fig. 3. Role of crucial amino acid residues of the CRAC sequence in agonist-stimulated calcium signaling of T2R4 receptors. Calcium mobilization in CHO cells expressing wild type (●, olive), Y114A (■, blue), and K117A (▲, maroon) receptors, is shown upon stimulation with increasing concentrations of quinine. All other conditions are as in Fig. 2. Data shown are means ± S.E. of at least three independent experiments. See [Materials and methods](#) for other details.




Fig. 4. Effect of membrane cholesterol content on agonist-stimulated calcium signaling in CHO cells expressing Y114A and K117A receptors. (a) Membrane cholesterol content of CHO cells transfected with Y114A (blue) and K117A (maroon) receptors upon cholesterol depletion and replenishment. Values are expressed as nmol of cholesterol per mg protein. Panels (b) and (c) show calcium signaling in control (●, olive), cholesterol-depleted (▲, blue), and cholesterol-replenished (■, maroon) CHO cells expressing (b) Y114A and (c) K117A receptors upon stimulation with increasing concentrations of quinine. All other conditions are as in Fig. 2. Data shown are means \pm S.E. of at least three independent experiments (* and *** correspond to significant ($p < 0.05$ and $p < 0.001$) difference in cholesterol content of cholesterol-depleted cells to control cells). See Materials and methods for other details.

3.5. Molecular docking of cholesterol to the T2R4 CRAC motif

In order to further understand the possible molecular interactions of the amino acid residues of the CRAC sequence of T2R4 and membrane cholesterol, we performed molecular docking of cholesterol with the CRAC motif. Cholesterol was docked to the CRAC motif of T2R4 (highlighted in Fig. 1), and the T2R4-cholesterol complex was energy minimized. Amino acid residues present within 4 Å of the cholesterol-binding pocket were analyzed for possible interactions. Cholesterol was docked toward the intracellular side of transmembrane helices III and IV. The polar hydroxyl group of cholesterol was pointed toward

the cellular interior and the flexible iso-octyl hydrocarbon tail facing the membrane. Fig. 5 shows the T2R4 CRAC motif residues that are involved in interaction with cholesterol. The nitrogen atom of Lys117 side chain was found to be involved in hydrogen bond interaction with the β -hydroxyl group of cholesterol. According to this model, Tyr114 does not interact with cholesterol, and its side chain appears to face transmembrane helices of T2R4. The hydrophobic region of cholesterol was flanked in the hydrophobic pocket formed by aliphatic amino acids Leu109, Ile112, Leu113, and Val116 of transmembrane helix III and Ile137 and Leu140 of transmembrane helix IV.

4. Discussion

As mentioned above, the role of membrane cholesterol in GPCR function constitutes an active and emerging area. Recent reports describing the presence of closely associated cholesterol molecules [31–34] in high resolution crystal structures of GPCRs have provided additional impetus in this area. In spite of a growing number of reports on cholesterol sensitivity of GPCR function, a general consensus on the mechanism of GPCR-cholesterol interaction remains elusive.

The human T2Rs belong to the GPCR superfamily. Their importance stems from the fact that a large number of structurally diverse and naturally occurring bitter compounds are recognized by them. Failure in detecting these compounds could be lethal since some of the bitter compounds are extremely toxic [10]. The perception of taste represents a crucial element because it acts as a sensor in the context of ingestion of food. Since cholesterol is an important part of human food, and cholesterol content in human tissues is contributed by food intake and metabolism [41], interaction of membrane cholesterol with bitter taste receptors could be relevant. In this work, we show that T2R4, a representative member of the bitter taste receptor family, exhibits sensitivity to membrane cholesterol in its function.




Fig. 5. Energy-minimized model of T2R4 receptor docked to cholesterol. Cholesterol was docked between the transmembrane helices III and IV in the CRAC motif of the receptor. Cholesterol is shown in pink and residues within 4 Å are shown in green. A hydrogen bond between K117 and β -OH group of cholesterol is shown as a broken blue line.

Membrane cholesterol has been shown to be crucial for the organization and function of a number of GPCRs [11,13,14,16]. As mentioned above, the effect of membrane cholesterol on GPCR organization and function could be due to direct and/or indirect effects [19]. In this context, a number of structural features of proteins have been suggested to be involved in preferential association with cholesterol [36,42]. Among these, one of the most studied motifs in membrane proteins, that exhibit sensitivity to cholesterol content, is the CRAC motif [35, 36]. Subsequent to their identification in peripheral-type benzodiazepine receptors [35], after more than a decade, CRAC motifs were identified in GPCRs such as rhodopsin, the β_2 -adrenergic receptor, the serotonin_{1A} receptor [37] and the human type I cannabinoid receptor [43]. In addition, membrane proteins such as caveolin-1 [44] and the HIV-1 transmembrane protein gp41 [45] have also been shown to possess CRAC motifs. In order to decipher the basis of the observed cholesterol sensitivity of T2R4 function, we examined whether bitter taste receptors in general, and T2R4 in particular, possess CRAC sequence. We report here the presence of CRAC motif in T2R4 and demonstrate that Lys117 plays an important role in its cholesterol sensitivity. To the best of our knowledge, our results represent the first report of the molecular basis of cholesterol sensitivity in the function of taste receptors. It is interesting to note that it has been recently proposed that cholesterol binding motif should contain a positively charged residue capable of participating in electrostatic interactions with 3 β -hydroxyl group of cholesterol [39,40]. Our present results with T2R4 therefore support this prediction and demonstrate that lysine in the CRAC sequence of T2R4 plays a crucial role in sensitivity of the receptor signaling to membrane cholesterol.

As stated above, the Y114A mutant did not exhibit any significant difference from the wild type receptor in terms of calcium mobilization (Fig. 3 and Table 1) and sensitivity to membrane cholesterol (Fig. 4b). Analysis of the energy-minimized model of T2R4 (Fig. 5) shows that the side chain of Tyr114 appears to face the transmembrane helices, away from the cholesterol. Our results therefore suggest that the Tyr114 does not appear to play a significant role in signaling by the receptor. However, we cannot exclude the possibility of either direct or indirect influence of other amino acids in T2R4-cholesterol interactions. It is important to note here that several amino acids at the interface of transmembrane helix III and intracellular loop 2, at which T2R4 has CRAC motif, have been shown to play a crucial role in G-protein coupling and signaling by GPCRs.

It is important to note here that the mere presence of CRAC motif(s) does not necessarily mean that this site preferentially associates with cholesterol. In other words, the presence of CRAC motif is only indicative of possible interaction of cholesterol with receptors and it would therefore be prudent to rely on further experimental validation of such interaction. For example, it has been shown that although the neurotensin receptor 1, a member of the GPCR family, possess two CRAC motifs, it does not display any dependence on membrane cholesterol for its function [20]. In this context, it is important to decipher molecular details of the nature of cholesterol sensitivity in case of bitter taste receptors, which would lead to better insight in our overall understanding of the functioning of these receptors in healthy and diseased states. Future analysis of the frequency of occurrence of mutant taste receptors in various populations could provide novel information on the development of gastronomic preferences across the world over the evolution of human civilization.

Conflict of interest

The authors declare no conflict of interest.

Transparency document

The Transparency document associated with this article can be found in the online version.

Acknowledgments

This work was supported by a Discovery grant (RGPIN-2014-04099) from the Natural Sciences and Engineering Research Council of Canada and a MMSF (# 8-2013-00) Allen Rouse Career Award (P.C.) and by the Council of Scientific and Industrial Research (Govt. of India) Network project BSC0115 (A.C.). S.P. was supported by a Graduate Studentship from Manitoba Health Research Council. A.C. gratefully acknowledges support from J.C. Bose Fellowship (Department of Science and Technology, Govt. of India). A.C. is an Adjunct Professor of Tata Institute of Fundamental Research (Mumbai), RMIT University (Melbourne, Australia), Indian Institute of Technology (Kanpur), and Indian Institute of Science Education and Research (Mohali). We thank G. Aditya Kumar for help in making figures, and members of the Chattopadhyay laboratory for their comments and discussions.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.bbamem.2016.06.005>.

References

- [1] K.L. Pierce, R.T. Premont, R.J. Lefkowitz, Seven-transmembrane receptors, *Nat. Rev. Mol. Cell Biol.* 3 (2002) 639–650.
- [2] D.M. Rosenbaum, S.G.F. Rasmussen, B.K. Kobilka, The structure and function of G-protein-coupled receptors, *Nature* 459 (2009) 356–363.
- [3] R. Heilker, M. Wolff, C.S. Tautermann, M. Bieler, G-protein-coupled receptor-focused drug discovery using a target class platform approach, *Drug Discov. Today* 14 (2009) 231–240.
- [4] A. Di Pizio, A. Levit, M. Slutzki, M. Behrens, R. Karaman, M.Y. Niv, Comparing class A GPCRs to bitter taste receptors: structural motifs, ligand interactions and agonist-to-antagonist ratios, *Methods Cell Biol.* 132 (2016) 401–427.
- [5] A. Jaggupilli, R. Howard, J.D. Upadhyaya, R.P. Bhullar, P. Chelikani, Bitter taste receptors: novel insights into the biochemistry and pharmacology, *Int. J. Biochem. Cell Biol.* (2016), <http://dx.doi.org/10.1016/j.biocel.2016.03.005> (in press).
- [6] J. Chandrashekar, K.L. Mueller, M.A. Hoon, E. Adler, L. Feng, W. Guo, C.S. Zuker, N.J.P. Ryba, T2Rs function as bitter taste receptors, *Cell* 100 (2000) 703–711.
- [7] N. Singh, S.P. Pydi, J. Upadhyaya, P. Chelikani, Structural basis of activation of bitter taste receptor T2R1 and comparison with class A G-protein-coupled receptors (GPCRs), *J. Biol. Chem.* 286 (2011) 36032–36041.
- [8] S.R. Foster, E. Roura, W.G. Thomas, Extrasensory perception: odorant and taste receptors beyond the nose and mouth, *Pharmacol. Ther.* 142 (2014) 41–61.
- [9] F.A. Shaik, N. Singh, M. Arakawa, K. Duan, R.P. Bhullar, P. Chelikani, Bitter taste receptors: extraoral roles in pathophysiology, *Int. J. Biochem. Cell Biol.* (2016), <http://dx.doi.org/10.1016/j.biocel.2016.03.011> (in press).
- [10] M. Behrens, W. Meyerhof, Bitter taste receptors and human bitter taste perception, *Cell. Mol. Life Sci.* 63 (2006) 1501–1509.
- [11] T.J. Pucadyil, A. Chattopadhyay, Role of cholesterol in the function and organization of G-protein coupled receptors, *Prog. Lipid Res.* 45 (2006) 295–333.
- [12] B. Chini, M. Parenti, G-protein-coupled receptors, cholesterol and palmitoylation: facts about fats, *J. Mol. Endocrinol.* 42 (2009) 371–379.
- [13] Y.D. Paila, A. Chattopadhyay, Membrane cholesterol in the function and organization of G-protein coupled receptors, *Subcell. Biochem.* 51 (2010) 439–466.
- [14] J. Oates, A. Watts, Uncovering the intimate relationship between lipids, cholesterol and GPCR activation, *Curr. Opin. Struct. Biol.* 21 (2011) 802–807.
- [15] A. Chattopadhyay, GPCRs: lipid-dependent membrane receptors that act as drug targets, *Adv. Biol.* 2014 (2014) 143023.
- [16] M. Jafurulla, A. Chattopadhyay, Membrane lipids in the function of serotonin and adrenergic receptors, *Curr. Med. Chem.* 20 (2013) 47–55.
- [17] M. Jafurulla, A. Chattopadhyay, Sphingolipids in the function of G protein-coupled receptors, *Eur. J. Pharmacol.* 763 (2015) 241–246.
- [18] D. Sengupta, A. Chattopadhyay, Molecular dynamics simulations of GPCR-cholesterol interaction: an emerging paradigm, *Biochim. Biophys. Acta* 1848 (2015) 1775–1782.
- [19] Y.D. Paila, A. Chattopadhyay, The function of G-protein coupled receptors and membrane cholesterol: specific or general interaction? *Glycoconj. J.* 26 (2009) 711–720.
- [20] J. Oates, B. Faust, H. Attrill, P. Harding, M. Orwick, A. Watts, The role of cholesterol on the activity and stability of neurotensin receptor 1, *Biochim. Biophys. Acta* 1818 (2012) 2228–2233.
- [21] S.P. Pydi, R.P. Bhullar, P. Chelikani, Constitutively active mutant gives novel insights into the mechanism of bitter taste receptor activation, *J. Neurochem.* 122 (2012) 537–544.
- [22] S.P. Pydi, N. Singh, J. Upadhyaya, R.P. Bhullar, P. Chelikani, The third intracellular loop plays a critical role in bitter taste receptor activation, *Biochim. Biophys. Acta* 1838 (2014) 231–236.
- [23] J. Upadhyaya, N. Singh, R.P. Bhullar, P. Chelikani, The structure-function role of C-terminus in human bitter taste receptor T2R4 signaling, *Biochim. Biophys. Acta* 1848 (2015) 1502–1508.

- [24] T.J. Pucadyil, A. Chattopadhyay, Cholesterol depletion induces dynamic confinement of the G-protein coupled serotonin_{1A} receptor in the plasma membrane of living cells, *Biochim. Biophys. Acta* 1768 (2007) 655–668.
- [25] S. Shrivastava, T.J. Pucadyil, Y.D. Paila, S. Ganguly, A. Chattopadhyay, Chronic cholesterol depletion using statin impairs the function and dynamics of human serotonin_{1A} receptors, *Biochemistry* 49 (2010) 5426–5435.
- [26] X. Li, L. Staszewski, H. Xu, K. Durick, M. Zoller, E. Adler, Human receptors for sweet and umami taste, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 4692–4696.
- [27] S. Kalipatnapu, T.J. Pucadyil, K.G. Harikumar, A. Chattopadhyay, Ligand binding characteristics of the human serotonin_{1A} receptor heterologously expressed in CHO cells, *Biosci. Rep.* 24 (2004) 101–115.
- [28] D.M. Amundson, M. Zhou, Fluorometric method for the enzymatic determination of cholesterol, *J. Biochem. Biophys. Methods* 38 (1999) 43–52.
- [29] R. Zidovetzki, I. Levitan, Use of cyclodextrins to manipulate plasma membrane cholesterol content: evidence, misconceptions and control strategies, *Biochim. Biophys. Acta* 1768 (2007) 1311–1324.
- [30] Y.D. Paila, E. Jindal, S.K. Goswami, A. Chattopadhyay, Cholesterol depletion enhances adrenergic signaling in cardiac myocytes, *Biochim. Biophys. Acta* 1808 (2011) 461–465.
- [31] V. Cherezov, D.M. Rosenbaum, M.A. Hanson, S.G.F. Rasmussen, F.S. Thian, T.S. Kobilka, H.-J. Choi, P. Kuhn, W.I. Weis, B.K. Kobilka, R.C. Stevens, High-resolution crystal structure of an engineered human β_2 -adrenergic G protein-coupled receptor, *Science* 318 (2007) 1258–1265.
- [32] M.A. Hanson, V. Cherezov, M.T. Griffith, C.B. Roth, V.-P. Jaakola, E.Y.T. Chien, J. Velasquez, P. Kuhn, R.C. Stevens, A specific cholesterol binding site is established by the 2.8 Å structure of the human β_2 -adrenergic receptor, *Structure* 16 (2008) 897–905.
- [33] W. Liu, E. Chun, A.A. Thompson, P. Chubukov, F. Xu, V. Katritch, G.W. Han, C.B. Roth, L.H. Heitman, A.P. Ijzerman, V. Cherezov, R.C. Stevens, Structural basis for allosteric regulation of GPCRs by sodium ions, *Science* 337 (2012) 232–236.
- [34] H. Wu, C. Wang, K.J. Gregory, G.W. Han, H.P. Cho, Y. Xia, C.M. Niswender, V. Katritch, J. Meiler, V. Cherezov, P.J. Conn, R.C. Stevens, Structure of a class C GPCR metabotropic glutamate receptor 1 bound to an allosteric modulator, *Science* 344 (2014) 58–64.
- [35] H. Li, V. Papadopoulos, Peripheral-type benzodiazepine receptor function in cholesterol transport. Identification of a putative cholesterol recognition/interaction amino acid sequence and consensus pattern, *Endocrinology* 139 (1998) 4991–4997.
- [36] R.M. Epand, Cholesterol and the interaction of proteins with membrane domains, *Prog. Lipid Res.* 45 (2006) 279–294.
- [37] M. Jafurulla, S. Tiwari, A. Chattopadhyay, Identification of cholesterol recognition amino acid consensus (CRAC) motif in G-protein coupled receptors, *Biochem. Biophys. Res. Commun.* 404 (2011) 569–573.
- [38] D. Sengupta, A. Chattopadhyay, Identification of cholesterol binding sites in the serotonin_{1A} receptor, *J. Phys. Chem. B* 116 (2012) 12991–12996.
- [39] N. Jamin, J.-M. Neumann, M.A. Ostuni, T.K.N. Vu, Z.-X. Yao, S. Murail, J.-C. Robert, C. Giatzakis, V. Papadopoulos, J.-J. Lacapère, Characterization of the cholesterol recognition amino acid consensus sequence of the peripheral-type benzodiazepine receptor, *Mol. Endocrinol.* 19 (2005) 588–594.
- [40] R.F. Epand, A. Thomas, R. Brasseur, S.A. Vishwanathan, E. Hunter, R.M. Epand, Juxtamembrane protein segments that contribute to recruitment of cholesterol into domains, *Biochemistry* 45 (2006) 6105–6114.
- [41] S.M. Grundy, Absorption and metabolism of dietary cholesterol, *Annu. Rev. Nutr.* 3 (1983) 71–96.
- [42] J. Fantini, F.J. Barrantes, How cholesterol interacts with membrane proteins: an exploration of cholesterol-binding sites including CRAC, CARC, and tilted domains, *Front. Physiol.* 4 (2013) 31.
- [43] S. Oddi, E. Dainese, F. Fezza, M. Lanuti, D. Barcaroli, V. De Laurenzi, D. Centonze, M. Maccarrone, Functional characterization of putative cholesterol binding sequence (CRAC) in human type-1 cannabinoid receptor, *J. Neurochem.* 116 (2011) 858–865.
- [44] R.M. Epand, B.G. Sayer, R.F. Epand, Caveolin scaffolding region and cholesterol-rich domains in membranes, *J. Mol. Biol.* 345 (2005) 339–350.
- [45] N. Vincent, C. Genin, E. Malvoisin, Identification of a conserved domain of the HIV-1 transmembrane protein gp41 which interacts with cholesterol groups, *Biochim. Biophys. Acta* 1567 (2002) 157–164.