

Effect of capsaicin on ligand binding activity of the hippocampal serotonin_{1A} receptor

Rajesh Prasad · Pushpendra Singh ·
Amitabha Chattopadhyay

Received: 31 May 2008 / Revised: 20 August 2008 / Accepted: 26 August 2008 / Published online: 19 September 2008
© Springer Science + Business Media, LLC 2008

Abstract The serotonin_{1A} receptor is an important member of the G-protein coupled receptor family, and is involved in the generation and modulation of a variety of cognitive, behavioral, and developmental functions. In order to examine the role of membrane material properties in ligand binding activity of the hippocampal serotonin_{1A} receptor, we monitored the function of the receptor in presence of capsaicin. Capsaicin has been previously shown to increase the elasticity of membrane bilayers. Our results show that the ligand binding activity of the hippocampal serotonin_{1A} receptor is reduced in the presence of capsaicin in a linear concentration-dependent manner. This is accompanied by no appreciable change in G-protein coupling of the receptor and overall membrane order. We conclude that material properties of membrane bilayers could play an important role in the function of the serotonin_{1A} receptor in particular, and membrane proteins in general.

Keywords Capsaicin · Hippocampal serotonin_{1A} receptor · Fluorescence anisotropy · Membrane material property · Ligand binding activity

Abbreviations

5-HT _{1A} receptor	5-hydroxytryptamine-1A receptor
DMPC	dimyristoyl- <i>sn</i> -glycero-3-phosphocholine
DMSO	dimethyl sulfoxide
DPH	1,6-diphenyl-1,3,5-hexatriene

GABA	γ -aminobutyric acid
GTP- γ -S	guanosine-5'- <i>O</i> -(3-thiotriphosphate)
GPCR	G-protein coupled receptor
PMSF	phenylmethylsulfonyl fluoride

Introduction

The seven transmembrane domain G-protein coupled receptor (GPCR) superfamily is one of the largest and most diverse protein families in mammals and constitute ~1–2% of the human genome [1]. The primary function of GPCRs is to transduce signals across membranes [2, 3]. GPCRs represent major targets for the development of novel drug candidates in all clinical areas [4]. The serotonin_{1A} (5-HT_{1A}) receptor is an important GPCR and is the most extensively studied among the serotonin receptors. The serotonin_{1A} receptor is involved in the generation and modulation of a variety of cognitive, behavioral, and developmental functions (for a recent review, see [5]). The serotonin_{1A} receptor agonists and antagonists represent major classes of molecules with potential therapeutic effects in anxiety- or stress-related disorders. Interestingly, mutant (knockout) mice lacking the serotonin_{1A} receptor exhibit enhanced anxiety-related behavior [6] and represent an important animal model for the analysis of complex traits such as anxiety disorders and aggression in higher animals [7, 8].

Since a significant portion of integral membrane proteins remains in contact with the membrane [9], the structure and function of membrane proteins often depend on the membrane lipid composition [10, 11]. The function of membrane proteins can be modulated either by specific (local) lipid–protein interactions, or material (physical)

R. Prasad · P. Singh · A. Chattopadhyay (✉)
Centre for Cellular and Molecular Biology,
Council of Scientific and Industrial Research,
Uppal Road, Hyderabad 500 007, India
e-mail: amit@ccmb.res.in

were treated with increasing concentrations of capsaicin for 1 h at 25°C, and 50 nmol of total phospholipid was suspended in 1.5 ml of buffer C and used for fluorescence anisotropy experiments. The amount of probe added was such that the final probe concentration was 1 mol% with respect to the total phospholipid content. This ensures optimal fluorescence intensity with negligible membrane perturbation. Membranes were vortexed for 1 min after addition of the probe and kept in the dark for 1 h before measurements. Background samples were prepared the same way except that the probe was omitted. The final probe concentration was 0.33 μM in all cases and the methanol content was low (~0.3%, v/v). Steady state fluorescence was measured in a Hitachi F-4010 spectrofluorometer using 1 cm path length quartz cuvettes at room temperature (~23°C). Excitation and emission wavelengths were set at 358 and 430 nm. Excitation and emission slits with nominal band passes of 1.5 and 20 nm were used. The excitation slit was kept low to avoid any photoisomerization of DPH. In addition, fluorescence was measured with a 30 s interval between successive openings of the excitation shutter (when the sample was in the dark in the fluorometer) to reverse any photoisomerization of DPH [32]. The optical density of the samples measured at 358 nm was 0.13±0.01. The anisotropy values remained identical even after dilution of membrane samples, indicating the absence of any scattering artifact [33]. Fluorescence anisotropy measurements were performed using a Hitachi polarization accessory. Anisotropy values were calculated from the equation [34]:

$$r = \frac{I_{VV} - GI_{VH}}{I_{VV} + 2GI_{VH}} \quad (1)$$

where I_{VV} and I_{VH} are the measured fluorescence intensities (after appropriate background subtraction) with the excitation polarizer vertically oriented and the emission polarizer vertically and horizontally oriented, respectively. G is the grating correction factor and is equal to I_{HV}/I_{HH} . All experiments were done with multiple sets of samples and average values of fluorescence anisotropy are shown in Fig. 3.

GTP-γ-S sensitivity assay In experiments with GTP-γ-S, agonist binding assays were carried out (as described above), with increasing concentrations of GTP-γ-S in the presence and absence of 0.25 mM capsaicin. The concentrations of GTP-γ-S leading to 50% inhibition of specific agonist binding (IC_{50}) were calculated by nonlinear regression fit of the data to a four parameter logistic function [35]:

$$B = [a/(1 + (x/I)^s)] + b \quad (2)$$

where B is the specific binding of the agonist normalized to agonist binding at the lowest concentration of GTP-γ-S, x

denotes concentration of GTP-γ-S, a is the range ($y_{\max} - y_{\min}$) of the fitted curve on the ordinate (y -axis), I is the IC_{50} concentration, b is the background of the fitted curve (y_{\min}) and s is the slope factor.

Statistical analysis Significance levels were estimated using student's two-tailed t -test using Graphpad Prism software version 4.00 (San Diego, CA, USA).

Results and discussion

Capsaicin has previously been reported to modulate function of membrane proteins [36]. Figure 2 shows the effect of increasing concentrations of capsaicin on specific [3 H]8-OH-DPAT binding to hippocampal serotonin $_{1A}$ receptors. The figure shows that agonist binding is reduced in a concentration-dependent manner with ~50% reduction in specific binding activity when 0.25 mM capsaicin was used. The agonist binding is reduced to ~5% of its original value when 0.5 mM capsaicin was used. We interpret the reduction in agonist binding activity in presence of capsaicin due to its ability to increase the elasticity of the membrane bilayer. It therefore appears that the effect of capsaicin on ligand binding activity is nonspecific, possibly




Fig. 2 Effect of capsaicin on specific binding of the agonist [3 H]8-OH-DPAT to the hippocampal serotonin $_{1A}$ receptor. Radioligand binding to the hippocampal serotonin $_{1A}$ receptor was monitored with increasing concentrations of capsaicin. A linear relationship of specific binding of [3 H]8-OH-DPAT to the hippocampal serotonin $_{1A}$ receptor as a function of increasing concentrations of capsaicin is observed (see inset). A linear regression analysis gave an excellent fit ($R^2 \sim 0.99$). Values are expressed as percentages of specific [3 H]8-OH-DPAT binding obtained in control membranes. Data shown are means±standard error from at least eight independent experiments in duplicate [*** correspond to p -value<0.0001, the differences in specific [3 H]8-OH-DPAT binding were tested against the corresponding value obtained in the absence of capsaicin (control value)]. See “Materials and methods” for other details




Fig. 3 Fluorescence anisotropy of the membrane probe DPH in hippocampal membranes monitored at varying concentrations of capsaicin. Fluorescence anisotropy measurements were carried out with membranes containing 50 nmol phospholipid at a probe to phospholipid ratio of 1:100 (mol/mol) at room temperature ($\sim 23^{\circ}\text{C}$). Data represent means \pm standard error from at least three independent experiments in duplicate. See “Materials and methods” for other details

due to change in elasticity of the membrane bilayer induced by capsaicin. This is supported by the linear relationship between the agonist binding activity and capsaicin concentration ($R^2 \sim 0.99$, see inset of Fig. 2).

In order to monitor the effect of capsaicin on membrane order, we carried out fluorescence anisotropy measurements with the membrane probe, DPH. Fluorescence anisotropy measured using probes such as DPH is correlated to its rotational diffusion [34], which is sensitive to the packing of lipid chains. This is due to the fact that fluorescence anisotropy depends on the degree to which the probe is able to reorient after excitation, and probe reorientation is a function of local lipid packing. DPH, which is a rod-like hydrophobic molecule, partitions into the interior (fatty acyl chain region) of the bilayer. Figure 3 shows that the fluorescence anisotropy of DPH exhibits a modest (up to $\sim 12\%$) increase with increasing concentrations of capsaicin. Similar change in fluorescence anisotropy of DPH [26] and other fluorescent probes [27] in presence of capsaicin have previously been reported.

As mentioned above, the reduction in agonist binding in presence of capsaicin could be attributed to its ability to increase the elasticity of the membrane bilayer rather than a direct effect. In such a scenario, it would be interesting to explore the effect of capsaicin on downstream signaling. To examine this, we monitored whether the G-protein coupling of the hippocampal serotonin_{1A} receptor is altered upon capsaicin treatment. Most of the seven transmembrane domain receptors are coupled to G-proteins [37], and guanine nucleotides are known to regulate agonist binding.

The serotonin_{1A} receptor is negatively coupled to adenylate cyclase system through G-proteins [38]. Sensitivity of agonist binding to guanine nucleotides can be monitored by performing the agonist binding assay in the presence of GTP- γ -S, a non-hydrolyzable analogue of GTP. We previously showed that the specific binding of the agonist [³H]8-OH-DPAT to hippocampal serotonin_{1A} receptors is sensitive to guanine nucleotides and is inhibited with increasing concentrations of GTP- γ -S. Figure 4 shows the inhibition of specific [³H]8-OH-DPAT binding to the serotonin_{1A} receptor in hippocampal membranes by GTP- γ -S in a characteristic concentration-dependent manner, with an estimated half maximal inhibition concentration (IC_{50}) of 77 nM. This shows that the hippocampal serotonin_{1A} receptor is coupled to G-proteins and GTP- γ -S induces a transition of the receptor from a high affinity to a low affinity state, as reported previously [39]. Interestingly, the inhibition curve in case of membranes treated with capsaicin (0.25 mM) does not display significant shift, as apparent from an estimated IC_{50} value of 70 nM. This shows that capsaicin treatment does not alter G-protein coupling of the serotonin_{1A} receptor.

Our results show that the ligand binding activity of the hippocampal serotonin_{1A} receptor is reduced in the presence of capsaicin, possibly due to increase in membrane elasticity (*i.e.*, decrease in membrane stiffness). Such alteration in ligand binding by membrane elasticity has previously been observed for the GABA_A receptor [21]. In addition, the function of membrane ion channels such as gramicidin [22], and the sodium channel [20] have been reported to be




Fig. 4 Effect of increasing concentration of GTP- γ -S on the specific binding of [³H]8-OH-DPAT to hippocampal serotonin_{1A} receptors in the presence (filled circle in solid line) or absence (empty circle in dashed line) of 0.25 mM capsaicin. Values are expressed as percentages of the specific binding obtained at the lowest concentration of GTP- γ -S. The curves are non-linear regression fit to the experimental data using Eq. 2. The data points represent means \pm standard error from at least five independent experiments in duplicate. See “Materials and methods” for other details

dependent on membrane elasticity. To the best of our knowledge, this constitutes the first report describing the role of membrane mechanical properties in ligand binding of this important neurotransmitter receptor (serotonin_{1A} receptor).

We have previously shown that depletion of membrane cholesterol results in reduction of the ligand binding activity of the hippocampal serotonin_{1A} receptor [12]. Interestingly, cholesterol depletion also increases bilayer elasticity (reduces stiffness) [22, 23]. Whether the loss in activity of the hippocampal serotonin_{1A} receptor, earlier reported by us [12], is due to a change in membrane elasticity induced by cholesterol depletion represents an interesting question. This could also be important in cases where the role of other types of membrane lipids (such as glycosphingolipids) are considered in relation to membrane protein function. We conclude that material properties of the bilayer are important in addressing the role of membrane lipids in the function of membrane proteins.

Acknowledgements This work was supported by the Council of Scientific and Industrial Research, Government of India. R.P. and P.S. thank the Council of Scientific and Industrial Research for the award of Research Fellowships. We gratefully acknowledge Dr. Olaf Andersen (Weill Medical College of Cornell University, NY, USA) for his kind gift of capsaicin for initial experiments. We thank Sourav Haldar for help with fluorescence measurements. A.C. is an Adjunct Professor at the Special Centre for Molecular Medicine of Jawaharlal Nehru University (New Delhi, India), and Honorary Professor of the Jawaharlal Nehru Centre for Advanced Scientific Research (Bangalore, India). We thank members of our laboratory for critically reading the manuscript.

References

1. Fredriksson, R., Schiöth, H.B.: The repertoire of G-protein-coupled receptors in fully sequenced genomes. *Mol. Pharmacol.* **67**, 1414–1425 (2005). doi:10.1124/mol.104.009001
2. Pierce, K.L., Premont, R.T., Lefkowitz, R.J.: Seven-transmembrane receptors. *Nat. Rev. Mol. Cell Biol.* **3**, 639–650 (2002). doi:10.1038/nrm908
3. Kroeze, W.K., Sheffler, D.J., Roth, B.L.: G-protein-coupled receptors at a glance. *J. Cell Sci.* **116**, 4867–4869 (2003). doi:10.1242/jcs.00902
4. Hopkins, A.L., Groom, C.R.: The druggable genome. *Nat. Rev. Drug Discov.* **1**, 727–730 (2002). doi:10.1038/nrd892
5. Pucadyil, T.J., Kalipatnapu, S., Chattopadhyay, A.: The serotonin_{1A} receptor: A representative member of the serotonin receptor family. *Cell. Mol. Neurobiol.* **25**, 553–580 (2005). doi:10.1007/s10571-005-3969-3
6. Julius, D.: Serotonin receptor knockouts: a moody subject. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 15153–15154 (1998). doi:10.1073/pnas.95.26.15153
7. Gingrich, J.A., Hen, R.: Dissecting the role of the serotonin system in neuropsychiatric disorders using knockout mice. *Psychopharmacology (Berl.)* **155**, 1–10 (2001). doi:10.1007/s002130000573
8. Toth, M.: 5-HT_{1A} receptor knockout mouse as a genetic model of anxiety. *Eur. J. Pharmacol.* **463**, 177–184 (2003). doi:10.1016/S0014-2999(03)01280-9
9. Lee, A.G.: Lipid-protein interactions in biological membranes: a structural perspective. *Biochim. Biophys. Acta.* **1612**, 1–40 (2003). doi:10.1016/S0005-2736(03)00056-7
10. Palsdottir, H., Hunte, C.: Lipids in membrane protein structures. *Biochim. Biophys. Acta.* **1666**, 2–18 (2004). doi:10.1016/j.bbamem.2004.06.012
11. Lee, A.G.: How lipids affect the activities of integral membrane proteins. *Biochim. Biophys. Acta.* **1666**, 62–87 (2004). doi:10.1016/j.bbamem.2004.05.012
12. Pucadyil, T.J., Chattopadhyay, A.: Cholesterol modulates the ligand binding and G-protein coupling to serotonin_{1A} receptors from bovine hippocampus. *Biochim. Biophys. Acta.* **1663**, 188–200 (2004). doi:10.1016/j.bbamem.2004.03.010
13. Paila, Y.D., Pucadyil, T.J., Chattopadhyay, A.: The cholesterol-complexing agent digitonin modulates ligand binding of the bovine hippocampal serotonin_{1A} receptor. *Mol. Membr. Biol.* **22**, 241–249 (2005). doi:10.1080/09687860500093453
14. Burger, K., Gimpl, G., Fahrenholz, F.: Regulation of receptor function by cholesterol. *Cell. Mol. Life Sci.* **57**, 1577–1592 (2000). doi:10.1007/PL00000643
15. Pucadyil, T.J., Chattopadhyay, A.: Role of cholesterol in the function and organization of G-protein coupled receptors. *Prog. Lipid Res.* **45**, 295–333 (2006). doi:10.1016/j.plipres.2006.02.002
16. Gimpl, G., Wiegand, V., Burger, K., Fahrenholz, F.: Cholesterol and steroid hormones: modulators of oxytocin receptor function. *Prog. Brain Res.* **139**, 43–55 (2002). doi:10.1016/S0079-6123(02)39006-X
17. Gimpl, G., Burger, K., Fahrenholz, F.: A closer look at the cholesterol sensor. *Trends Biochem. Sci.* **27**, 596–599 (2002). doi:10.1016/S0968-0004(02)02224-7
18. Ohvo-Rekilä, H., Ramstedt, B., Leppimäki, P., Slotte, J.P.: Cholesterol interactions with phospholipids in membranes. *Prog. Lipid Res.* **41**, 66–97 (2002). doi:10.1016/S0163-7827(01)00020-0
19. Grinthal, A., Guidotti, G.: Bilayer mechanical properties regulate the transmembrane helix mobility and enzymatic state of CD39. *Biochemistry.* **46**, 279–290 (2007). doi:10.1021/bi061052p
20. Lundbaek, J.A., Birn, P., Hansen, A.J., Søgaard, R., Nielsen, C., Grishman, J., et al.: Regulation of sodium channel function by bilayer elasticity: the importance of hydrophobic coupling. Effects of micelle-forming amphiphiles and cholesterol. *J. Gen. Physiol.* **123**, 599–621 (2004). doi:10.1085/jgp.200308996
21. Søgaard, R., Werge, T.M., Bertelsen, C., Lundbye, C., Madsen, K. L., Nielsen, C.H., et al.: GABA_A receptor function is regulated by lipid bilayer elasticity. *Biochemistry.* **45**, 13118–13129 (2006). doi:10.1021/bi060734+
22. Lundbaek, J.A., Birn, P., Grishman, J., Hansen, A.J., Andersen, O. S.: Membrane stiffness and channel function. *Biochemistry.* **35**, 3825–3830 (1996). doi:10.1021/bi952250b
23. Needham, D., Nunn, R.S.: Elastic deformation and failure of lipid bilayer membranes containing cholesterol. *Biophys. J.* **58**, 997–1009 (1990)
24. Kuenzi, F.M., Dale, N.: Effect of capsaicin and analogues on potassium and calcium currents and vanilloid receptors in *Xenopus* embryo spinal neurones. *Br. J. Pharmacol.* **119**, 81–90 (1996)
25. Lundbaek, J.A., Birn, P., Tape, S.E., Toombes, G.E.S., Søgaard, R., Koeppe II, R.E., et al.: Capsaicin regulates voltage-dependent sodium channels by altering lipid bilayer elasticity. *Mol. Pharmacol.* **68**, 680–689 (2005)
26. Meddings, J.B., Hogaboam, C.M., Tran, K., Reynolds, J.D., Wallace, J.L.: Capsaicin effects on non-neuronal plasma membranes. *Biochim. Biophys. Acta.* **1070**, 43–50 (1991). doi:10.1016/0005-2736(91)90144-W
27. Aranda, F.J., Villalain, J., Gómez-Fernández, J.C.: Capsaicin affects the structure and phase organization of phospholipids membranes. *Biochim. Biophys. Acta.* **1234**, 225–234 (1995). doi:10.1016/0005-2736(94)00293-X

28. Tsuchiya, H.: Biphasic membrane effects of capsaicin, an active component in *Capsicum* species. *J. Ethnopharmacol.* **75**, 295–299 (2001). doi:[10.1016/S0378-8741\(01\)00200-8](https://doi.org/10.1016/S0378-8741(01)00200-8)
29. Smith, P.K., Krohn, R.I., Hermanson, G.T., Mallia, A.K., Gartner, F.H., Provenzano, M.D., et al.: Measurement of protein using bicinchoninic acid. *Anal. Biochem.* **150**, 76–85 (1985). doi:[10.1016/0003-2697\(85\)90442-7](https://doi.org/10.1016/0003-2697(85)90442-7)
30. Bruns, R.F., Lawson-Wendling, K., Pugsley, T.A.: A rapid filtration assay for soluble receptors using polyethylenimine-treated filters. *Anal. Biochem.* **132**, 74–81 (1983). doi:[10.1016/0003-2697\(83\)90427-X](https://doi.org/10.1016/0003-2697(83)90427-X)
31. McClare, C.W.F.: An accurate and convenient organic phosphorus assay. *Anal. Biochem.* **39**, 527–530 (1971). doi:[10.1016/0003-2697\(71\)90443-X](https://doi.org/10.1016/0003-2697(71)90443-X)
32. Chattopadhyay, A., London, E.: Fluorimetric determination of critical micelle concentration avoiding interference from detergent charge. *Anal. Biochem.* **139**, 408–412 (1984). doi:[10.1016/0003-2697\(84\)90026-5](https://doi.org/10.1016/0003-2697(84)90026-5)
33. Lentz, B.R., Moore, B.M., Barrow, D.A.: Light-scattering effects in the measurement of membrane microviscosity with diphenylhexatriene. *Biophys. J.* **25**, 489–494 (1979)
34. Lakowicz, J.R.: Principles of fluorescence spectroscopy, 3rd edn. Springer, New York (2006)
35. Higashijima, T., Ferguson, K.M., Sternweis, P.C., Smigel, M.D., Gilman, A.G.: Effects of Mg^{2+} and the beta gamma-subunit complex on the interactions of guanine nucleotides with G proteins. *J. Biol. Chem.* **262**, 762–766 (1987)
36. Rizvi, S.I., Luqman, S.: Capsaicin-induced activation of erythrocyte membrane sodium/potassium and calcium adenosine triphosphatases. *Cell. Mol. Biol. Lett.* **8**, 919–925 (2003)
37. Clapham, D.E.: The G-protein nanomachine. *Nature.* **379**, 297–299 (1996). doi:[10.1038/379297a0](https://doi.org/10.1038/379297a0)
38. Emerit, M.B., El Mestikawy, S., Gozlan, H., Rouot, B., Hamon, M.: Physical evidence of the coupling of solubilized 5-HT_{1A} binding sites with G regulatory proteins. *Biochem. Pharmacol.* **39**, 7–18 (1990). doi:[10.1016/0006-2952\(90\)90642-X](https://doi.org/10.1016/0006-2952(90)90642-X)
39. Harikumar, K.G., Chattopadhyay, A.: Differential discrimination of G-protein coupling of serotonin_{1A} receptors from bovine hippocampus by an agonist and an antagonist. *FEBS Lett.* **457**, 389–392 (1999). doi:[10.1016/S0014-5793\(99\)01082-0](https://doi.org/10.1016/S0014-5793(99)01082-0)